

FRiSCy Oxford Fracture Prevention Service

MK Javaid

Lecturer in Metabolic Bone Disease, University of Oxford

Hon Consultant Rheumatologist, Nuffield Orthopaedic Centre

Lead Clinician for FRiSCy

Declaration

These views are my own

- In last five years received honoraria, travel and/or subsistence expenses from:
 - Amgen, Eli Lilly, Medtronic, Norvartis, Proctor and Gamble, Servier, Shire

Mapping to Public Health

Responsibilities

- making the public healthier by encouraging discussions, advising government and supporting action by local government, the NHS and other people and organisations
- supporting the public so they can protect and improve their own health
- protecting the nation's health through the national health protection service, and preparing for public health emergencies
- sharing our information and expertise with local authorities, industry and the NHS, to help them make improvements in the public's health
- researching, collecting and analysing data to improve our understanding of health and come up with answers to public health problems
- reporting on improvements in the public's health so everyone can understand the challenge and the next steps
- helping local authorities and the NHS to develop the public health system and its specialist workforce

Mapping to Public Health

Priorities

- helping people to live longer and more healthy lives by reducing preventable deaths and the burden of ill health associated with smoking, high blood pressure, obesity, poor diet, poor mental health, insufficient exercise, and alcohol
- reducing the burden of disease and disability in life by focusing on preventing and recovering from the conditions with the greatest impact, including dementia, anxiety, depression and drug dependency

Lifetime fracture risk in 50 white men and women

	Men (%)	Women (%)
Hip	10.7	22.9
Clinical vertebral	8.3	15.1
Distal forearm	4.6	20.8
Humerus	4.1	12.9
Any of above	22.4	46.4

Lifetime risk for Wrist/Hip or vertebral is 30-40% (similar to CHD)

After a HIP fracture:

More bed days than heart attack and stroke combined

1:10 dead 1 month

1:3 dead by 12 months

50% require care home admission

2.3 billion in hip fracture costs alone per year

£2.3 billion / years more bed days than heart attack and stroke combined

340,00 fragility fractures last year

- > 68,000 Hip fractures last year
- > Each hip fracture costs £30K/2 years (NHFD 2010)




£2.3 billion / years more bed days than heart attack and stroke combined

- 340,00 fragility fractures last year
- > 68,000 Hip fractures last year
- > Each hip fracture costs £30K/2 years (NHFD 2010)


50% have already had a fracture

Drugs cost from £11 per year

RCT demonstrate reduced risk of re-fracture by 20 – 70%




DXA scan costs £69...



Primary or secondary prevention

Limited resource: maximize value






50% hip fracture patients have already had a fragility fracture

A fragility fracture Doubles risk of fracture


"After the first fracture, act to prevent the second"

Less than 50% of fracture patients are assessed for osteoporosis




Less than 50% of patients who start treatment are still on it at 1 year...


Achieving the outcome



High value




Delivered within Primary care: QOF 2012/13




MISSED treatment in 129,880- 146,534 (72 - 85%) of eligible patients

The NHS pathway



Already paying for finding and treating Hip fracture best practice tariff Un-provided other inpatient fracture (50%) & all outpatient fracture patients

The Effective Service



Minimally disruption healthcare

Public health, primary or secondary care led but *must* be patient focused

A therapy for every patient (annual cost)

ORAL

Alendronate 70 mg weekly (£11.44)
Risedronate 35mg weekly (£15.21)
Ibandronate 150 mg monthly (£18.98)
Strontium 2g nocte (£353)

Calcium replete

Vitamin D replete

Zoledronate 5mg iv annually (day case rate)

Denosumab 60mg sc 6m (£366)

Treatment adherence – options with parenteral therapy

Teriparatide 20mcg od s.c (£3263)

Fracture Reduction in South Central Policy group



A network of every bone clinician/ Nurse



AIM:

Every patient with a fragility fracture over the age of 50yr in South Central is:

- Identified
- Assessed
- 3. Treated effectively for at least five years for both bone and falls health


Fracture Liaison Service > Fracture Prevention Service


POSITIVE LIBERTY- We choose the scope of what we can decide






- Minimize patient burden
- Interface service
 - Trauma case finding + community monitoring
- Transfer care from high cost specialist settings to lower cost community settings
- Integrated care
 - Cost effective...
 - Safety
 - Patient Experience






Secondary fracture assessments by month




Prescribing of Bisphosphonates and other drugs affecting bone metabolism


Osteoporosis Prescribing Costs per Weighted Population (Cost per Astro PU)
October to December 2009



Fracture Prevention Specialist Nurse Recommendation (n=3,680)



Outcome of monitoring



Mapping to public health

Responsibilities

- making the public healthier by encouraging discussions, advising government and supporting action by local government, the NHS and other people and organisations
- supporting the public so they can protect and improve their own health
- protecting the nation's health through the national health protection service, and preparing for public health emergencies
- sharing our information and expertise with local authorities, industry and the NHS, to help them make improvements in the public's health
- researching, collecting and analysing data to improve our understanding of health and come up with answers to public health problems
- reporting on improvements in the public's health so everyone can understand the challenge and the next steps
- helping local authorities and the NHS to develop the public health system and its specialist workforce

Mapping to public health

Priorities

- helping people to live longer and more healthy lives by reducing preventable deaths and the burden of ill health associated with smoking, high blood pressure, obesity, poor diet, poor mental health, insufficient exercise, and alcohol
- reducing the burden of disease and disability in life by focusing on preventing and recovering from the conditions with the greatest impact, including dementia, anxiety, depression and drug dependency

Set the outcome: need to ensure all fragility fracture patients over 50 years have 4 steps





Find them

Assess them

Decide which treatment

Make sure they stay on it





BOTTOM

Champion network: Work out how much and how to do this for the locality

Summary



- Set the standards and priorities
- Population approach across 4 steps
- High value interventions
- Bone health and falls to reduce fracture









Kerri Rance

Sally Hope

Sarah James

Terri Morgan

Carol Weeks

Rachael Knight

Tracy Dobbin

Rhys Davies

Suzanne Jones/ Sylvie Thorn

Adam Dawroski / Antoinette Broad

Professors

Cooper, Wass, Willett, Arden, Carr

D Prieto Alhambra, A Judge, R Batra, G Round, A Kiran, K Leyland, A Soni, R Warne

And all the FRiSCy team!
Stephen, Colin, Annie, Gill, Mark...

ElevenInformatics:

John, Naveed & Joanna





